




PATIENT & COMMUNITY EDUCATION SERIES

Understanding Liver Tumors

A plain-language, comprehensive guide to liver tumors — the common benign lesions found incidentally on scans, how they're distinguished from cancer, and how each type is managed.

Written for anyone told "we found something on your liver" during an unrelated scan

Health Education Series · EDU-LIV

A publication of The Liver and Kidneys Hope Foundation · a Nevada 501(c)(3) nonprofit

First Edition

Important This guide is provided for general educational purposes only. The Liver and Kidneys Hope Foundation is a nonprofit focused on liver-health education, community support, and outreach — it is **not a medical provider** and does not offer diagnosis, treatment, or medical advice. Nothing here should replace a conversation with a qualified, licensed healthcare professional who knows your history.

CONTENTS

What's Inside

Hearing "there's a spot on your liver" after a scan done for something completely unrelated can be frightening — but the large majority of liver tumors found this way are entirely benign, and many need nothing more than confirmation and, at most, simple monitoring. This guide walks through what these findings actually are, and how doctors tell them apart.

1	Your Liver: A Quiet, Hardworking Organ	>
2	What Is a Liver Tumor? Understanding the Term	>
3	The Most Common Benign Tumor: Hemangioma	>
4	Focal Nodular Hyperplasia (FNH)	>
5	Hepatic Adenoma	>
6	Liver Cysts: Simple & Complex	>
7	Other Benign Liver Lesions	>
8	Malignant Liver Tumors: A Brief Overview	>
9	How Common Are Liver Tumors?	>
10	Signs & Symptoms	>
11	The Incidental Finding: What Happens Next	>
12	Imaging: Telling Tumors Apart	>
13	When Biopsy Is (and Isn't) Needed	>
14	Risk Factors for Different Tumor Types	>
15	Possible Complications of Benign Tumors	>
16	Treatment & Management: The Big Picture	>
17	Watchful Waiting: When Monitoring Is Right	>
18	When Treatment Is Needed: Procedures & Surgery	>



- 19 Hepatic Adenoma & Hormonal Considerations >
- 20 Living with a Diagnosed Liver Tumor >
- 21 Prevention & Myths vs. Facts >
- ★ Questions, Glossary, Resources & Disclaimer >

A NOTE BEFORE YOU BEGIN

This guide focuses primarily on **benign** liver tumors — the kind found incidentally, which make up the large majority of liver lesions discovered this way. For a full, dedicated look at malignant (cancerous) liver tumors — hepatocellular carcinoma, cholangiocarcinoma, and cancer that has spread to the liver — our companion "Understanding Liver Cancer" guide covers that territory in depth; Section 8 here gives a brief overview and points you there.

SECTION 1 Your Liver: A Quiet, Hardworking Organ

Before discussing what a "spot" on the liver might mean, it helps to appreciate just how much your liver does every day — usually without you noticing at all.

Your liver is the largest solid organ inside your body, weighing roughly 1.4 kg (about 3 pounds) in an adult. It sits just under your rib cage on the right side, tucked below the diaphragm and above the stomach and intestines. Scientists have identified more than **500 distinct functions** it performs, and its rich, dual blood supply (from both the hepatic artery and the portal vein) makes it a common organ to be closely examined on abdominal imaging done for almost any reason.

Why liver findings are so often incidental

Because the liver sits within view on so many routine scans — an ultrasound for gallstones, a CT scan after an injury, an MRI for back pain — it's one of the organs most likely to have something noticed on it purely by chance, unrelated to why the scan was originally ordered. This is precisely the scenario this guide is built around.

DID YOU KNOW?

Advances in imaging technology over the past few decades mean scans today detect far smaller, more subtle liver findings than were ever visible before — part of why incidental liver "spots" are reported more often now than in the past, without necessarily reflecting any actual increase in disease.

KEY POINT

Finding something on the liver during a scan for an unrelated reason is common, and the large majority of the time, it turns out to be one of the harmless, well-understood findings covered throughout this guide.

SECTION 2

What Is a Liver Tumor? Understanding the Term

"Tumor" is a word that simply means an abnormal growth or mass of tissue — it says nothing on its own about whether that growth is harmful.

Medically, a tumor is just a mass — a collection of cells growing somewhat differently from surrounding tissue. Tumors are broadly divided into two categories:

- **Benign tumors** — growths that don't invade surrounding tissue or spread elsewhere in the body. The large majority of liver tumors found incidentally fall into this category.
- **Malignant tumors (cancer)** — growths capable of invading nearby tissue and spreading to other parts of the body, covered briefly in Section 8 and in depth in our companion liver cancer guide.

Why "tumor" causes so much unnecessary alarm

Because "tumor" is so often used interchangeably with "cancer" in everyday language, hearing the word attached to your own liver can feel alarming before you even know which category applies. Understanding this basic distinction — that "tumor" is a neutral, descriptive term, not a diagnosis of cancer — is one of the most calming, useful pieces of information in this entire guide.

KEY POINT

When a clinician says "we found a liver tumor" or "a liver lesion," this is a description of an imaging finding, not a diagnosis — the entire rest of this guide is essentially about how that finding gets sorted into a specific, understood category.

DID YOU KNOW?

Radiologists often use the word "lesion" rather than "tumor" specifically to avoid this unintended alarm — the two terms are generally used interchangeably to mean "an abnormal area seen on imaging," without implying anything about whether it's benign or malignant.

SECTION 3

The Most Common Benign Tumor: Hemangioma

Hemangiomas are the single most common benign liver tumor, made up of a tangle of blood vessels — and the large majority never cause any problems at all.

What a hemangioma is

A hemangioma is a benign growth composed of a cluster of abnormal blood vessels, essentially present from birth, though most are only discovered later in life, incidentally, on imaging done for another reason. They are not cancerous, do not become cancerous, and generally don't require any treatment.

How they're typically identified

Hemangiomas have a distinctive appearance on contrast-enhanced imaging (particularly a specific pattern of how contrast fills the lesion over time), which often allows confident diagnosis through imaging alone, without needing a biopsy.

Do they need treatment?

The overwhelming majority of hemangiomas, including surprisingly large ones, cause no symptoms and require no treatment — simply confirming the diagnosis is often the entire extent of "management" needed. Very large hemangiomas occasionally cause symptoms like abdominal discomfort or a feeling of fullness, and rarely, treatment may be considered for a symptomatic or unusually large lesion.

KEY POINT

If you've been told you have a liver hemangioma, this is generally reassuring news — it's the most common benign liver finding, doesn't turn into cancer, and rarely requires anything beyond confirming the diagnosis.

DID YOU KNOW?

Hemangiomas are found in a meaningful share of the general population on imaging done for unrelated reasons, making them one of the most frequently encountered "incidental findings" in all of abdominal imaging.

SECTION 4

Focal Nodular Hyperplasia (FNH)

Focal nodular hyperplasia, often shortened to FNH, is the second most common benign liver tumor — a distinctive growth thought to arise from a localized area of increased blood flow.

What FNH is

FNH is a benign mass made up of normal liver cells arranged in an abnormal pattern, generally believed to develop in response to a small, localized abnormality in blood vessels. It is not cancerous and does not become cancerous.

Who it typically affects

FNH is considerably more common in women than men, and is often identified in young to middle-aged adults, frequently as an incidental finding.

A distinctive imaging appearance

FNH often has a specific feature on imaging — a central scar, visible with certain imaging techniques — that can help confidently distinguish it from other liver lesions, sometimes avoiding the need for a biopsy.

Do they need treatment?

Like hemangiomas, FNH rarely causes symptoms and generally requires no treatment once confidently diagnosed — simple confirmation and, in some cases, periodic monitoring is typically all that's needed.

KEY POINT

FNH is another example of a common, entirely benign finding that, once properly identified, usually requires little more than reassurance and, at most, occasional follow-up imaging.

DID YOU KNOW?

Unlike hepatic adenoma (Section 5), FNH does not appear to be strongly linked to hormonal contraceptive use, though some specialists still recommend periodic monitoring in women taking these medications out of general caution.

SECTION 5 Hepatic Adenoma

Hepatic adenoma is a benign liver tumor that deserves particular attention, because unlike hemangioma and FNH, it carries a small but real risk of bleeding and, in some cases, of transforming into cancer over time.

What a hepatic adenoma is

An adenoma is a benign growth of liver cells that, unlike FNH, isn't simply a normal-pattern overgrowth — it has its own somewhat abnormal cellular characteristics, which is part of why it's managed somewhat more cautiously than the other benign lesions covered so far.

The hormonal connection

Hepatic adenomas are strongly linked to estrogen exposure, occurring considerably more often in women who use, or have used, hormonal contraceptives (particularly older, higher-dose formulations) for extended periods. They can also occur, less commonly, in men, and in association with anabolic steroid use.

Why they're managed more cautiously

Larger adenomas carry a meaningful risk of spontaneous bleeding, which can occasionally be significant. A smaller subset of adenomas, particularly certain specific subtypes identifiable

through specialized imaging or biopsy, and larger adenomas, carry a small but real risk of transforming into liver cancer over time — the reasons management decisions (Section 18) are more individualized for this specific type of benign tumor.

KEY POINT

A hepatic adenoma diagnosis, unlike hemangioma or FNH, generally warrants a more active conversation about size, hormonal exposure, and monitoring or treatment — this is the one benign liver tumor type in this guide where "just reassurance" isn't usually the complete answer.

DID YOU KNOW?

Stopping hormonal contraceptives after a hepatic adenoma diagnosis can, in some cases, lead to the adenoma shrinking over time — a genuinely actionable, non-invasive first step your care team may recommend, discussed further in Section 19.

SECTION 6

Liver Cysts: Simple & Complex

Liver cysts — fluid-filled sacs within the liver — are extremely common incidental findings, and the large majority are entirely simple and harmless.

Simple cysts

A simple liver cyst is a thin-walled sac containing clear fluid, generally present from birth (congenital) and often not discovered until incidentally found later in life. Simple cysts are extremely common, entirely benign, and essentially never require treatment unless they become unusually large and symptomatic.

Complex cysts

Occasionally, a cyst has features on imaging that don't fit the simple pattern — internal septations (dividing walls), thickened walls, or solid components. These "complex" cysts warrant closer evaluation, since a small minority can represent something other than a simple cyst, including, rarely, infection, a parasitic cyst, or, uncommonly, a cystic tumor.

Polycystic liver disease

Some people have numerous liver cysts, sometimes associated with a genetic condition also affecting the kidneys (polycystic kidney disease). This pattern is generally still benign but may warrant its own specific monitoring approach given the larger number of cysts involved.

KEY POINT

A "simple" cyst designation on your imaging report is genuinely reassuring language — it means the finding fits a well-recognized, harmless pattern that essentially never requires any treatment or even follow-up imaging.

DID YOU KNOW?

Simple liver cysts are found in a meaningful share of the general population on abdominal imaging, making them one of the single most common incidental liver findings of all.

SECTION 7 Other Benign Liver Lesions

Beyond hemangioma, FNH, adenoma, and cysts, a handful of other benign findings are worth knowing about, even though they're encountered less frequently.

Focal fatty change (or focal fat sparing)

In people with fatty liver disease (MASLD, covered in our companion guide), fat can sometimes accumulate — or, conversely, be notably absent — in a specific, localized area of the liver, creating an appearance on imaging that can look like a distinct mass. This is not a true tumor at all, and specific imaging techniques can usually confirm this pattern confidently.

Regenerative nodules

In people with cirrhosis, areas of regenerating liver tissue can form nodules that may appear on imaging and require careful distinction from early liver cancer, covered in our companion cirrhosis and liver cancer guides.

Nodular regenerative hyperplasia

A less common, more diffuse pattern of small regenerative nodules throughout the liver, sometimes associated with certain other medical conditions or medications, generally managed with monitoring.

KEY POINT

Because several benign conditions can mimic a true tumor on imaging, radiologists and hepatologists rely on specific imaging patterns, and sometimes follow-up imaging or biopsy, to confidently sort a finding into the right category.

DID YOU KNOW?

Focal fatty change is becoming an increasingly common source of incidental liver findings, tracking the rising prevalence of MASLD — a good example of how one common condition can create imaging findings that need careful, specific interpretation.

SECTION 8

Malignant Liver Tumors: A Brief Overview

While this guide focuses on benign findings, it's worth briefly placing malignant liver tumors in context, since distinguishing the two is exactly what the diagnostic process described later in this guide is designed to do.

Primary liver cancer

Cancer that starts in the liver itself — most commonly hepatocellular carcinoma (HCC), which develops overwhelmingly in people with existing cirrhosis, and less commonly cholangiocarcinoma, arising from bile duct cells. Our companion liver cancer guide covers both in full depth, including staging, treatment, and surveillance.

Secondary (metastatic) liver tumors

Cancer that started elsewhere in the body (commonly the colon, breast, lung, or pancreas) and has spread to the liver. These are treated according to the original cancer's own guidelines, often alongside liver-directed treatments.

Why this distinction matters for an incidental finding

Unlike the benign lesions covered in this guide, malignant tumors generally require prompt, active treatment. This is precisely why the evaluation process described in Sections 11–13 exists — to reliably sort an incidental finding into the correct category, since benign and malignant tumors require completely different next steps.

KEY POINT

If your evaluation raises any concern for a malignant tumor, our companion "Understanding Liver Cancer" guide is the right, more detailed resource — this guide's role is helping you understand the broader landscape of possibilities and the process of sorting them out.

DID YOU KNOW?

The large majority of incidentally found liver lesions in someone without cirrhosis or a known cancer history turn out to be benign — risk factors like existing cirrhosis or a history of cancer elsewhere in the body meaningfully shift how a new finding is approached, covered further in Section 14.

SECTION 9 How Common Are Liver Tumors?

Benign liver tumors and lesions are remarkably common — likely affecting a substantial share of the general population, even though most people never know it.

Population studies using sensitive imaging have found simple cysts, hemangiomas, and other benign findings in a meaningful proportion of people undergoing abdominal imaging for entirely unrelated reasons — reflecting how common these findings are in people who have no liver disease and no symptoms whatsoever.

Why detection has increased

As mentioned in Section 1, improvements in imaging technology and the sheer increase in how often abdominal imaging is performed for various reasons have both increased how often these incidental findings are detected and reported — without necessarily reflecting any real increase in how common these growths actually are.

DID YOU KNOW?

This pattern — increasingly sensitive technology finding more incidental abnormalities that turn out to be harmless — is sometimes discussed under the broader concept of "incidentaloma," a term used across many areas of medicine, not just the liver, to describe this exact phenomenon.

KEY POINT

If you've been told about an incidental liver finding, you are genuinely far from alone — this is one of the most common types of incidental findings in all of medical imaging.

SECTION 10 Signs & Symptoms

The overwhelming majority of benign liver tumors cause no symptoms whatsoever — symptoms, when they do occur, generally relate to size rather than the specific type of tumor.

When symptoms can occur

- **Abdominal discomfort or a feeling of fullness**, generally only with unusually large lesions.
- **A palpable mass**, in rare cases of very large tumors.
- **Sudden abdominal pain**, which can indicate bleeding into a tumor (most relevant for hepatic adenoma, Section 5) and warrants prompt medical evaluation.

Why most liver tumors are found incidentally, not through symptoms

Because the liver has substantial spare capacity and few pain-sensing nerves, even a reasonably sized benign tumor typically doesn't affect overall liver function or cause noticeable symptoms — which is exactly why the overwhelming majority are found on imaging done for a completely unrelated reason.

SEEK MEDICAL CARE PROMPTLY

Sudden, severe abdominal pain, particularly with a known hepatic adenoma or other liver tumor, can indicate bleeding and warrants prompt medical evaluation.

KEY POINT

The absence of symptoms tells you very little about what type of liver finding you have — nearly all benign liver tumors, and even many early cancers, cause no symptoms at all, which is why imaging characteristics (Section 12), not how you feel, drive the diagnostic process.

SECTION 11

The Incidental Finding: What Happens Next

Understanding the typical pathway that follows an incidental liver finding can genuinely ease some of the uncertainty of waiting for answers.

Step 1 — The initial finding

A scan done for an unrelated reason (a CT for abdominal pain, an ultrasound for gallstones, an MRI for back pain) notes an abnormal area in the liver, generally described by size, location, and some basic characteristics.

Step 2 — Dedicated liver imaging

Often, the next step is a dedicated, more detailed scan of the liver specifically — typically a contrast-enhanced CT or MRI — designed to characterize the finding more precisely than the original scan, which wasn't necessarily optimized for evaluating the liver.

Step 3 — Comparison with any prior imaging

If you've had any previous abdominal imaging, your radiologist will often want to compare the new finding against old scans — a lesion that's been stable in size and appearance for years is generally far more reassuring than one seen for the first time.

Step 4 — Further evaluation, if needed

Depending on what the dedicated imaging shows, this may be the end of the process (with a confident, reassuring diagnosis), or it may lead to a recommendation for follow-up imaging in a few months, referral to a hepatologist, or, less commonly, a biopsy (Section 13).

KEY POINT

This staged process — starting broad, then getting progressively more specific — is designed to reach a confident answer as efficiently as possible, using the least invasive approach that will reliably do so.

TALK WITH YOUR CLINICIAN

It's entirely reasonable to ask directly: "What's the plan for figuring out what this finding is, and how long will that take?"

SECTION 12 Imaging: Telling Tumors Apart

Modern imaging can distinguish between many types of liver tumors with real confidence, often without ever needing a biopsy — understanding how helps explain why imaging plays such a central role.

Why contrast-enhanced imaging matters so much

Different types of liver tumors have their own blood supply patterns, and different tissue characteristics that behave differently when contrast dye is injected during a CT or MRI scan. Radiologists look at exactly how a lesion "lights up" and fades on serial images taken over time (called enhancement pattern) — this pattern is often distinctive enough to confidently identify the specific type of tumor present.

Tumor type	General imaging clue
Hemangioma	A characteristic, gradual, peripheral-to-central filling pattern.
FNH	Rapid, uniform enhancement, often with a visible central scar.
Simple cyst	No enhancement at all, with imaging characteristics consistent with simple fluid.
Hepatocellular carcinoma	A specific pattern of rapid uptake followed by rapid washout of contrast, discussed in our companion liver cancer guide.

MRI vs. CT

MRI often provides more detailed tissue characterization than CT for certain liver lesions, and doesn't involve radiation exposure, though CT remains excellent for many purposes and is more widely available and faster to perform — your radiologist and clinician will recommend whichever is most appropriate for your specific situation.

KEY POINT

For many benign liver tumors, a well-performed, dedicated contrast study can provide enough confidence to avoid a biopsy altogether — imaging alone is often genuinely sufficient.

DID YOU KNOW?

Because these enhancement patterns are so well-studied and specific, radiologists specializing in liver imaging can often provide a confident diagnosis directly in their report, sometimes phrased as "imaging findings are classic for [specific benign lesion]."

It's worth understanding why this level of confidence is possible for the liver specifically, more so than for many other organs. The liver's dual blood supply, combined with decades of research correlating specific enhancement patterns with confirmed pathology from biopsies and surgical specimens, has built an unusually detailed, reliable imaging playbook. This is part of why liver lesion characterization by imaging alone has become such a well-established, trusted practice in radiology.

SECTION 13

When Biopsy Is (and Isn't) Needed

Because imaging alone often confidently identifies many benign liver tumors, biopsy is used more selectively for liver lesions than for many other suspected tumors elsewhere in the body.

When biopsy is often avoided

When a lesion has classic, unambiguous imaging characteristics for a benign finding like hemangioma, FNH, or a simple cyst, biopsy is generally unnecessary and, in some cases, actively avoided — particularly for hemangiomas, where biopsy carries a small bleeding risk without adding meaningful diagnostic value once imaging is confident.

When biopsy may be recommended

- **When imaging findings are ambiguous** or don't clearly fit a specific benign pattern.
- **When distinguishing a hepatic adenoma from other lesions**, or characterizing its specific subtype, would change management.
- **When malignancy cannot be confidently excluded** by imaging alone.
- **When a person has significant risk factors**, such as underlying cirrhosis or a known cancer elsewhere in the body, which lowers the threshold for tissue confirmation.

What a liver biopsy involves

A thin needle, usually guided by imaging, removes a small tissue sample, typically under local anesthesia. Most people go home the same day, with some soreness at the biopsy site.

KEY POINT

If your care team recommends against a biopsy despite an unfamiliar-sounding finding, this often reflects genuine confidence from imaging alone, not an oversight — it's entirely reasonable to ask directly why biopsy isn't being recommended, if you're curious.

TALK WITH YOUR CLINICIAN

Worth asking: "Are you confident enough from imaging alone, or would a biopsy add useful information in my specific case?"

SECTION 14 Risk Factors for Different Tumor Types

Certain personal factors shift the likelihood of different tumor types, and factor directly into how an incidental finding is interpreted and managed.

Factor	Relevance
Existing cirrhosis	Substantially raises the likelihood that a new liver lesion could represent cancer, changing the diagnostic approach considerably (covered in our companion liver cancer guide).
History of cancer elsewhere in the body	Raises the possibility that a new liver lesion could represent metastatic spread.
Hormonal contraceptive or anabolic steroid use	A specific risk factor for hepatic adenoma.
Female sex	Associated with higher rates of both FNH and hepatic adenoma.
Underlying MASLD/fatty liver	Can produce focal fatty change or sparing that mimics a true tumor on imaging.

KEY POINT

The same imaging finding can be interpreted quite differently depending on these personal risk factors — which is why an accurate, complete medical history is just as important as the imaging itself in reaching the right diagnosis.

DID YOU KNOW?

Because cirrhosis so significantly changes the interpretation of any new liver finding, people with known cirrhosis are generally already undergoing regular surveillance imaging (covered in our companion cirrhosis and liver cancer guides), meaning new findings are often caught within this existing monitoring framework rather than as a first-time incidental discovery.

SECTION 15

Possible Complications of Benign Tumors

While benign liver tumors are, by definition, not cancerous, a small number can still cause specific complications worth understanding.

Bleeding

Larger hepatic adenomas carry the most meaningful bleeding risk among the benign tumors covered in this guide, occasionally causing sudden, significant internal bleeding — the reason size and growth are actively monitored for this specific tumor type (Section 18).

Malignant transformation

As covered in Section 5, certain hepatic adenoma subtypes and larger adenomas carry a small but real risk of transforming into liver cancer over time — a risk essentially not shared by hemangioma, FNH, or simple cysts, which do not undergo malignant transformation.

Symptoms from size alone

Very large benign tumors of any type can occasionally cause discomfort or a feeling of fullness simply by taking up space, or, rarely, by pressing on nearby structures.

Rupture during pregnancy

Hepatic adenomas can grow during pregnancy due to hormonal changes, with an increased risk of bleeding in this specific setting — a consideration covered further in Section 19.

KEY POINT

Nearly all of the meaningful complication risk in this section relates specifically to hepatic adenoma — hemangioma, FNH, and simple cysts carry essentially none of these risks once properly diagnosed.

DID YOU KNOW?

This concentration of risk in one specific tumor type is exactly why hepatic adenoma receives its own more active monitoring and treatment conversation, while the other benign tumors in this guide are generally managed with simple reassurance.

SECTION 16

Treatment & Management: The Big Picture

Management of benign liver tumors ranges from simple reassurance to active monitoring to, in select cases, treatment — depending entirely on the specific tumor type, size, and individual circumstances.

The general approach by tumor type

Tumor type	Typical management
Hemangioma	Reassurance; treatment only rarely, for very large or symptomatic lesions.
FNH	Reassurance, with occasional follow-up imaging.
Simple cyst	Reassurance; essentially never requires treatment.
Hepatic adenoma	Individualized — may include hormonal changes, monitoring, or procedural treatment, depending on size and features (Sections 18–19).
Complex cyst or ambiguous finding	Further evaluation, sometimes including biopsy, to reach a confident diagnosis before deciding on management.

A WORD OF CAUTION

The right management approach depends on your specific tumor type, size, and individual risk factors. **This guide cannot recommend a specific management plan for you.** These decisions belong with your treating clinician.

KEY POINT

For the majority of benign liver tumors, "management" genuinely means simple reassurance and, at most, periodic monitoring — not every finding requires active treatment, and knowing this can meaningfully ease the anxiety of an unfamiliar diagnosis.

SECTION 17

Watchful Waiting: When Monitoring Is Right

For many benign liver tumors, periodic monitoring — rather than immediate treatment — is the appropriate, evidence-based approach.

Why monitoring, rather than treatment, is often right

Because most benign liver tumors carry very low complication risk, and because any treatment (even a minimally invasive one) carries its own risks, monitoring a stable, low-risk lesion over time is often the more sensible approach than intervening on something unlikely to ever cause a problem.

What monitoring typically involves

Follow-up imaging at defined intervals (commonly a repeat scan at 6 to 12 months, sometimes longer if stability is confirmed) to confirm the lesion isn't growing or changing in a

concerning way. For hepatic adenoma specifically, this monitoring is often somewhat closer, given the complication risks covered in Section 15.

What "stable" means, and why it's reassuring

A lesion that looks the same on repeat imaging, months or years apart, is behaving in a way entirely consistent with a benign process — malignant tumors, by contrast, generally show growth or change over comparatively short periods of time.

KEY POINT

Being told "we'll just keep an eye on it" is a legitimate, evidence-based medical decision, not a sign that your clinician is unsure or dismissive — for many benign liver tumors, this genuinely is the best available approach.

TALK WITH YOUR CLINICIAN

If you're on a monitoring plan, it's worth clarifying: "How often will we repeat imaging, and for how long, before we can consider this fully resolved?"

It can help to think of watchful waiting as an active, deliberate strategy rather than a passive default — your care team has weighed the specific characteristics of your finding against the risks of intervention and concluded that observation offers the best balance for your situation. This is the same evidence-based reasoning used across many areas of medicine when a finding is low-risk and a treatment carries risks of its own; it reflects confidence in the diagnosis, not uncertainty about what to do next.

SECTION 18

When Treatment Is Needed: Procedures & Surgery

For the minority of benign liver tumors that do require active treatment — most often larger or symptomatic hepatic adenomas — several effective options exist.

Surgical removal (resection)

Surgically removing the tumor, leaving the remaining healthy liver in place, is generally considered for larger hepatic adenomas (commonly those above a certain size threshold), certain adenoma subtypes with higher malignant transformation risk, or any tumor causing significant symptoms.

Minimally invasive procedures

For select smaller lesions, ablation techniques (similar to those used for early liver cancer, covered in our companion liver cancer guide) or embolization procedures may be considered as an alternative to open surgery.

Management of hormonal factors first

For hepatic adenoma specifically, stopping hormonal contraceptives is often tried first, particularly for smaller lesions, since this alone can lead to shrinkage in some cases, potentially avoiding the need for a procedure altogether (covered further in Section 19).

KEY POINT

Treatment, when needed, is generally reserved for the specific situations where the benefit clearly outweighs the risk of the procedure itself — not a default response to any benign tumor diagnosis.

A WORD OF CAUTION

Whether treatment is appropriate, and which type, depends on your specific tumor characteristics. **This guide cannot recommend a specific procedure for you.** These decisions belong with your treating specialist.

SECTION 19

Hepatic Adenoma & Hormonal Considerations

Because hepatic adenoma is so closely tied to hormone exposure, this specific tumor type deserves its own dedicated look at contraception and pregnancy considerations.

Hormonal contraceptives

If you're diagnosed with a hepatic adenoma while using hormonal contraceptives (particularly combined oral contraceptives), your clinician will very likely discuss stopping them, since this alone can lead to stabilization or shrinkage of the adenoma in many cases. Alternative, non-hormonal contraceptive options can be discussed with your care team.

Pregnancy

Pregnancy causes significant hormonal changes that can cause existing hepatic adenomas to grow, with an increased risk of bleeding during pregnancy specifically. If you have a known hepatic adenoma and are pregnant or planning a pregnancy, this warrants proactive discussion with both your hepatologist and obstetric team, ideally before conception where possible, so a monitoring or treatment plan can be established in advance.

Men and anabolic steroids

While less common than the female, contraceptive-related pattern, hepatic adenoma can occur in men, particularly with anabolic steroid use — stopping the causative substance is generally the first recommended step in this situation as well.

KEY POINT

Because hormonal factors so directly influence hepatic adenoma behavior, addressing the hormonal exposure itself is often the single most impactful, least invasive first step in management.

TALK WITH YOUR CLINICIAN

If you have a hepatic adenoma and are on hormonal contraception, or are pregnant or planning to be, raise this proactively — timing and coordination between specialists genuinely matters here.

SECTION 20 Living with a Diagnosed Liver Tumor

For the large majority of people with a benign liver tumor, daily life is entirely unaffected once a confident diagnosis is reached — this section covers the emotional side of getting there.

The anxiety of waiting for answers

The period between an initial incidental finding and a confident, final diagnosis can be genuinely stressful, even when the eventual answer turns out to be reassuring. This anxiety is completely understandable, and it's reasonable to ask your care team for a realistic timeline and to check in if that timeline stretches longer than expected.

After a reassuring diagnosis

Once a benign diagnosis like hemangioma, FNH, or a simple cyst is confirmed, most people can set the finding aside entirely, without ongoing worry or, in many cases, even further monitoring — a genuinely reassuring place to land after the uncertainty of the workup.

Living with an adenoma under monitoring

For those with a hepatic adenoma under active monitoring, periodic imaging appointments are a manageable, routine part of life for most people — a small, worthwhile inconvenience given the protective value of catching any change early.



IF YOU ARE STRUGGLING

If you are experiencing persistent low mood, hopelessness, or thoughts of harming yourself, please reach out to a healthcare professional or a crisis line. The **988 Suicide & Crisis Lifeline** (call or text 988 in the United States) is free and available 24/7. Support is available.

COMMUNITY MATTERS

If you're navigating an unusually complex or ongoing liver tumor evaluation, organizations like The Liver and Kidneys Hope Foundation can offer general support and connection while you work through the diagnostic process with your care team.

SECTION 21 Prevention & Myths vs. Facts

Prevention here looks a little different than for many other conditions in this guide series, since most benign liver tumors aren't preventable in the usual sense — but a few genuinely relevant points are worth knowing.

Prevention

- **Being mindful of long-term hormonal contraceptive or anabolic steroid use**, given the specific link to hepatic adenoma.
- **Managing MASLD/fatty liver**, which can reduce the confusing focal fatty changes that sometimes mimic a true tumor on imaging.
- **Following recommended surveillance** if you have cirrhosis, so any new finding is caught within an established monitoring framework.

Myths vs. facts

Myth	Fact
"A liver tumor always means cancer."	"Tumor" simply means an abnormal growth — the large majority found incidentally are entirely benign.
"If they're not doing a biopsy, they must be missing something."	Imaging alone often confidently diagnoses many benign liver tumors, making biopsy genuinely unnecessary in many cases.
"All benign liver tumors need to be removed eventually."	Most benign liver tumors, including hemangioma, FNH, and simple cysts, never require any treatment.
"If I have a liver cyst, it will definitely grow and cause problems."	Simple cysts are typically stable over time and essentially never require treatment.
"Watchful waiting means my doctor isn't taking this seriously."	Monitoring is a legitimate, evidence-based approach for many benign liver tumors, chosen because it's genuinely the most appropriate option.
"Once I've had one incidental liver finding, I'll probably keep finding more."	Having one benign lesion doesn't meaningfully increase your likelihood of developing others — most people with a confirmed benign finding never have another incidental discovery.
"A 'lesion' on my report sounds worse than a 'tumor'."	Radiologists use these terms interchangeably to mean an abnormal area on imaging — neither word implies anything about whether it's benign or malignant.

KEY POINT

If you remember one thing from this page: an incidental liver finding is common, the word "tumor" doesn't mean cancer, and the diagnostic process described in this guide is specifically designed to reach a confident, reassuring answer for the large majority of people.

CLOSING

Questions, Glossary, Resources & Disclaimer

Questions you might ask your doctor

- What specific type of liver finding is this, based on imaging?
- How confident are you in this diagnosis from imaging alone, or would a biopsy add useful information?
- Do I need any follow-up imaging, and if so, when and how often?

- If this is a hepatic adenoma, should I reconsider my hormonal contraception?
- What symptoms, if any, should prompt me to seek care sooner?
- At what point can we consider this fully resolved, without further monitoring?

Glossary

Term	Plain-language meaning
Benign	Not cancerous; doesn't invade nearby tissue or spread elsewhere.
Enhancement pattern	How a lesion "lights up" and fades on contrast imaging, used to help identify its type.
FNH (focal nodular hyperplasia)	A common benign liver tumor made of normal liver cells in an abnormal pattern.
Hemangioma	The most common benign liver tumor, made of a tangle of blood vessels.
Hepatic adenoma	A benign liver tumor linked to hormone exposure, with a small risk of bleeding or malignant transformation.
Incidentaloma	An abnormal finding discovered incidentally on imaging done for an unrelated reason.
Lesion	A general term for an abnormal area seen on imaging, not implying benign or malignant.

About The Liver and Kidneys Hope Foundation

The Liver and Kidneys Hope Foundation (TLHF) is a Nevada-based 501(c)(3) nonprofit dedicated to liver-health education, community support, and outreach. Through educational materials like this guide, community programming, our chapter network, volunteer efforts, and fundraising initiatives, we work to raise awareness of liver conditions and support the people affected by them, including those navigating an incidental liver tumor finding. The Foundation is not a medical provider and does not deliver clinical care.

A NOTE ON SOURCES AND CURRENCY

This guide reflects widely accepted, general medical understanding of benign liver tumors intended for patient education. It does not cite individual studies. For the most current, individualized information about your specific finding, your healthcare team is the best resource.



Full disclaimer

PLEASE READ

This publication is provided by The Liver and Kidneys Hope Foundation for **educational purposes only**. It is **not medical advice** and is not a substitute for professional diagnosis, treatment, or the judgment of a qualified, licensed healthcare provider who knows your personal history. The Liver and Kidneys Hope Foundation is **not a medical provider** and does not practice medicine, provide clinical services, or establish a provider-patient relationship through this material.

Do not use this guide to diagnose or treat a health problem or disease, and do not disregard or delay seeking professional medical advice because of something you have read here. Sudden, severe abdominal pain can indicate a medical emergency — contact your local emergency services immediately. If you think you may have a medical emergency for any other reason, do the same.

The Foundation makes no warranties about the completeness, accuracy, or currency of this information and accepts no liability for actions taken based on it. Always consult your own healthcare professional.